

EXAMINATION OF ENDOCRINE DISORDERS (2)

Case reports

Prof. Dr. Pavel Maruna

Case report No. 1
Hypothyroidism



Female, 46 yr.

Patient treated with arterial hypertension for last 3 years;

Palpitations for many years, established diagnosis of neuro-circulatory asthenia, higher frequency of palpitations in last months, reported as paroxysmal heart pulsation, accompanied with a pant and shortness of breath, independent on her physical activity;

Impaired glucose tolerance, last 3 yr. on a diet;

As a new problem, paresthesia in fingers, swelling of hands developed several weeks ago.

Allergy: negative

Chronic medication: ramipril (angiotensin-converting enzyme inhibitor) as an anti-hypertensive treatment

Abuse: alcohol and smoking - negative

Gynecological history: menses from 12 to 44 yr. (e.g., menopause 2 yr. ago), 2 deliveries, 1 abortion.

Social situation: married, 2 kids, working as a shop assistant.

Family history: twin sister treated with thyroid dysfunction; without other serious diseases in her family.

Actual problems:

Progressive difficulties last year:

Patient is slow, non effective, sleepy.

Tiredness, worsening of memory.

Poorer pronunciation, rhonchus (=low pitch chest wheezing).

Constipation.

Weight increase about 5 kg last 12 months.

Physical examination:

**Blood pressure 130 / 85 mmHg, heart rate 50 / min., regular,
temperature 36,0 °C**

Height 165 cm, weight 74 kg, BMI 27,2

Differences from a normal status:

- **Moderated/ slowed down psycho-motor tempo,**
- **limited mimics in face,**
- **rhonchus,**
- **dry skin with a yellow tone,**
- **spare hair,**
- **symmetrically edematous (upper) eyelids,**
- **symmetric soft painless goiter,**
- **hand and finger swelling,**
- **peri-maleolar ankle edemas.**

Laboratory findings

Na ⁺	138 mmol/l	(137 - 146)
K ⁺	4,1 mmol/l	(3,8 - 5,0)
Cl ⁻	101 mmol/l	(97 - 108)
glycaemia	5,5 mmol/l	(3,9 - 5,6)
urea	6,9 mmol/l	(2,8 - 8,0)
creatinine	112 µmol/l	(44 - 110)
cholesterol	7,8 mmol/l	(3,8 - 5,8)
triacyl-glycerols	1,1 mmol/l	(0,68 - 1,69)
total bilirubin	17 mmol/l	(2,0 - 17,0)
ALT	0,46 ukat/l	(0,10 - 0,78)
AST	0,72 ukat/l	(0,10 - 0,72)

Blood count

Leukocytes	8,6 x 10 ³ /mm ³	(4,1 - 10,2)
Erythrocytes	3,12 x 10 ⁶ /mm ³	(4,19 - 5,75)
Hemoglobin	101 g/l	(135 - 174)
HCT	32,8	(0,39 - 0,51)
MCV	105 fl	(82 - 98)
MCH	0,32 pg	(28,0 - 34,6)
Platelets	295 x 10 ³ /mm ³	(142 - 327)

Thyroid gland examinations

free T4	4,2 pmol/l	(11,0 - 21,0)
free T3	0,7 nmol/l	(1,5 - 3,0)
TSH	33,3 mIU/l	(0,5 - 5,0)
anti-TSHR Ab	negative	
anti-TPO Ab	+++	
anti-thyroglobulin Ab	++	

USG imaging

Volume of thyroid gland 25 ml (normal limits to 20 ml), lobes are symmetric, bilaterally numerous foci with low echo-genicity. Thyroid gland delimited (without invasion to surrounding structures). Lymph nodes of normal shape.

Achilles tendon reflex:

markedly protracted

ECG

Sinus rhythm, ventricular activity 50 / min., horizontal axis, PR segment 220 ms, QRS duration 90 ms, QT-c interval 420 ms, low voltage of QRS complex, ST segment in iso-electric line, normal shape of T wave.

Recommendation:

To start a treatment with levothyroxine (Euthyrox tbl.) in a dose 25 ug per day (1 tbl. in the morning).

After 4 weeks, Euthyrox may be increased to 50 ug per day.

After 2 months, laboratory control of TSH, free T3 and free T4 levels.

Question No. 1

Diagnosis of hypothyroidism in this patient can be established with a knowledge of history and physical examination, still before laboratory examination.

Nevertheless, which laboratory parameters confirm hypothyroidism in this patient? What is their relation to thyroid dysfunction ?

Question No. 2

**Mild anemia was found in this patients.
Based on morphological classification, which type of anemia is it?**

What is a pathogenesis of anemia in hypothyroidism ?

Question No. 3

Discuss following findings and their possible links to hypothyroidism:

- **hypercholesterolemia**
- **ECG curve changes**

Question No. 4

The diagnosis of thyroid disorder is based on knowledge of:

- 1. Thyroid function (hypothyroidism, hyperthyroidism, euthyroid status);**
- 2. Local signs (goiter, local expansion - trachea, esophagus, vessels);**
- 3. Etiological classification.**

What is a cause (etiology) of hypothyroidism in this patient?



Case report No. 2

Osteomalacia



History

Recorded in 2008

Female, 60 yr. old

Past history: without severe diseases in anamnesis

Chronic treatment: irregularly analgetic drugs for dorsalgia (back pain)

Allergy: negative

Abuse: negative

Family history: without important diseases in parents and siblings

Gyn. history: delivery 1, abortion 0.

Social status: lives in solitude, old-age pensioner

History

Patient was admitted to internal department of regional hospital in Moravia for progressive dyspnoe stage II.-III. (according to NYHA classification)

In physical examination

cachexia,

heavy kypho-scoliosis,

short stature (136 cm),

pectus excavatum

tachypnoe, dyspnoe in minimal activities (walking).

In laboratory – **markedly elevated D- dimers (>6000)**,

(norm is less than 500 nanograms per milliliter, ng/mL)

high activity of **ALP 16 μ kat / l** (norm is less than 2.2 μ kat / l).

History

The examination in hospital excluded following diseases:

**acute coronary syndrome,
pulmonary embolism,
dissection of aortal aneurysma,
pneumotorax**

...

X-ray imaging

Chest native X-ray: Heavy kyphoscoliosis of thoracic backbone.

Backbone skiagram: Diffuse osteoporosis with de-differentiation and destruction of vertebrae.

Skull skiagram: Without destructive changes of bones.

**With these findings, the patient was send to hospitalization
to Osteological department of the 3rd Internal clinic,
1st Faculty of Medicine in Prague
...to examine bone metabolism and to start a therapy.**

**With these findings, the patient was send to hospitalization
to Osteological department of the 3rd Internal clinic,
1st Faculty of Medicine in Prague
...to examine bone metabolism and to start a therapy.**

**Newly patient reports a progressive lowering in body
height in last 2 years.**

As a main problem, patient complains about dorsal pain.

Physical examination

Height **136 cm**, weight **29 kg**, BMI **16.1**

BP 130/90 mmHg, heart rate 115/min.,

O₂ saturation 95%

Kypho-scoliosis, pectus excavatum.

Physical examination

Height **136 cm**, weight **29 kg**, BMI **16.1**

BP 130/90 mmHg, heart rate 115/min.,

O₂ saturation 95%

Kyphoscoliosis, pectus excavatum.

ECG

Sinus tachycardia 105 / min., AV blockade of the 1st degree

Laboratory examination

Na ⁺	141 mmol/l	(137 - 146)
K ⁺	5,0 mmol/l	(3,8 - 5,0)
Cl ⁻	107 mmol/l	(7 - 108)
Ca ²⁺	2,66 mmol/l	(2,00 - 2,65)
Ca ²⁺	1,45 mmol/l	(1,0 - 1,4)
H ₂ PO ₄ ⁻	0,93 mmol/l	(0,6 - 1,6)
Mg	0,79 mmol/l	(0,66 - 1,08)
LDH	2,63 ukat/l	(2,20 - 3,75)
Urea	6,0 mmol/l	(2,8 - 8,0)
Creatinine	61,0 µmol/l	(44,0 - 110,0)
Uric acid	411 mmol/l	(220 - 420)

Laboratory examination

ALT	0,26 µkat/l	(0,10 - 0,78)
AST	0,28 µkat/l	(0,10 - 0,72)
GGT	0,66 µkat/l	(0,14 - 0,84)
Bilirubin	8 mmol/l	(2,0 - 17,0)
ALP	13,53 µkat/l	(0,6 - 2,2)
Cholesterol	5,4 mmol/l	(3,8 - 5,8)
Triacylglycerols	1,8 mmol/l	(0,68 - 1,69)
Total protein	78 g/l	(65,0 - 85,0)
Albumin	40 g/l	(35,0 - 53,0)
TSH	6,821 mIU/l	(0,5 - 5,0)
Free T4	15.2 pmol/l	(11,0 - 21,0)
PTH	257,2 pmol/l	(1,6 - 6,9)

USG of thyroid and parathyroid glands

Enlargement of a parathyroid body bellow left lobe of thyroid gland.
Thyroid gland of a normal size and shape.


^{99m}Tc -MIBI

Higher accumulation of ^{99m}Tc -MIBI near a lower edge of a left thyroid lobe

99mTc-MIBI

= methoxy isobuthyl isonitril

The molecule passes cells membranes passively, once intracellular it further accumulates in the mitochondrias. Detection of ^{99m}Tc gamma emission



Adenoma of a parathyroid gland



Abdominal USG imaging

Without pathological changes.

Echocardiography

Normal ventricular function, normal valvular morphology and function.

Spirometry

Severe restrictive respiratory disease.

Severe reduction of a vital lung capacity to 25% of normal values.


Diagnostic consideration (1)

Findings of \uparrow PTH, \uparrow Ca, and HPO_4 near a lower limit support a diagnosis of **primary hyperparathyroidism**.
A severe bone syndrome in a clinical and laboratory examination.

Suspect **adenoma** of a left lower parathyroid gland.

There is an indication to surgical removal of parathyroid adenoma.

...And following surgical treatment of deforming kyphoscoliosis with markedly decreased lung capacity.




X-ray examination of pelvis and femoral bones

Marked deformation of pelvic skeleton, „heart-shaped“ pelvis, deflection of both femoral bones, low density of all displayed bones

Chest X-ray image

**Bell-shaped chest with gibbus,
Total resorption of humerus head
on the left side.**




Chest X-ray image

Bell-shaped chest with gibbus,
Total resorption of humerus head
on the left side.

Conclusion

Skeletal deformities are typical
for **osteomalacia**



„Bone“ laboratory parameters


Osteocalcin	201,8 µg/l	(15 - 46)
25(OH)-D₃ vitamin	5 ng/ml	(15 - 42)
1,25(OH)-D₃ vitamin	48,4 ng/l	(19 - 67)
U - Ca	2,5 mmol/24 h	(3,5 - 6,5)
U - HPO₄	19,1 mmol/24 h	(13 - 42)

Diagnostic consideration (2)


Skeletal deformities are characteristic for **osteomalacia**, not for primary hyperparathyroidism.

Extremely low levels of **calcidiol** (25-OH-D₃ vitamin), as a marker of vitamin D storage in organism.

7-Dehydrocholesterol (liver)



Photoreaction
(skin)




Calcitriol
(cholecalciferol, D₃)

25-hydroxylation (liver)
1 α -hydroxylation (kidneys)

Vitamin D₃ metabolism

Calcitriol
(1 α ,25-dihydroxycalcitriol)



Diagnostic consideration (2)

Skeletal deformities are characteristic for **osteomalacia**, not for primary hyperparathyroidism.

Extremely low levels of **kalcidiol** (25-OH-D₃ vitamin), as a marker of vitamin D storage in organism.

Marked skeletal decalcification – low **calciuria**, **high ALP** (marker of increased osteoblastic activity).

Diagnostic consideration (2)


Findings give evidence against primary hyperparathyroidism

Long-term hypo-calcemia leads to hyperplasia of parathyroid glands with following autonomous over-function of one of them.

This course suggests for **tertiary hyperparathyroidism**.

Tertiary hyperparathyroidism

Due to prolonged hypocalcemia, one hyperplastic parathyroid body developed functionally autonomous adenoma. Hypocalcemia has changed to hypercalcemia.



Recommendation

- **Surgical revision of parathyroid glands**, displacement of autonomous body (v. s. adenoma) as well as other 3 (likely hyperplastic) parathyroid bodies, the maintenance of $\frac{1}{2}$ of one of them - what is sufficient for normal PTH production.
- Post-surgery massive **re-calcification** due to expected „hungry bone“ syndrome.

Surgery

February 20 – 29, 2008, hospitalization at the 3rd Dept. of surgery, Faculty hospital in Motol.

February 22, 2008 – Extirpation of adenoma of the left lower parathyroid gland, currently total extirpation of upper and lower parathyroid body on the right side, subtotal ablation of the left upper body.

Post-surgical course:

Laboratory: Ca 1,71....1,75 mmol / l

Post-surgery creepiness of hands and face ... as expected manifestation of transitory hypocalcemia

2nd hospitalization

February 29 – March 02, 2008, 3rd Internal Department

Subjectively: Intermittently depressive.

Objectively: Rest scar above sternum, without other variations.

Laboratory:


- Ca 1,85 mmol/l (2,0 - 2,65)
- HPO₄ 0,67 mmol/l (0,6 - 1,6)
- ALP 25,45 ukat/l (0,6 - 2,2)
- PTH 18,01 pmol/l (1,6 - 6,9).

Diagnostic consideration (3)

1. **Tertiary hyperparathyroidism** – extirpation of parathyroid adenoma which developed in long-term secondary hyperparathyroidism.
2. **Post-surgery „hungry bone“** bone syndrome with hypocalcemia.
3. **Osteomalacia** of an unclear origin as a primary disease

Ca balance in organism

Daily intake
approx. 25 - 30 mmol
(phosphates, oxalates...)



Absorption
upto 15 mmol / d

Secretion
to 7 mmol / d

17 - 25 mmol / d

ECT


2,5 mmol / l
totally only 35 mmol

ICT

signal function
0,1 μ mol / l

Filtration
240 mmol/d

Tubular resorption
230 mmol / d




5 - 6 mmol / d

Bone remodeling

7,5 mmol / d

Quick replacement
500 mmol / d

Mineral deposits
25 mol Ca
(rapidly replaceable
only 100 mmol)



What now?

1. To establish etiology of osteomalacia – suspicion of malabsorption.
2. Repeated controls of calcemia during therapy:
Calcium 1,5 g / den, Alpha-D3 1 ug/day.

(Alpha-D3 is vitamin D3 hydroxylated in position 1)

Hospitalization on the 4th Internal Clinic, 1st Faculty of Medicine in Prague

**Diagnostic hospitalization for suspicion of intestinal
malabsorption syndrome**

Auto-Ab, coeliac disease

Auto-antibodies negative

Gastroscopy

Normal macroscopic finding

Biopsy with normal histological image

Colonoscopy

Diverticulis of sigmoideum, other parts without pathological changes.

Histological examination from biopsies with normal findings.



Abdominal CT imaging

Without important changes. As an incidental finding, a cyst of right hepatic lobe and a renal stone of a right kidney (10x7 mm) were revealed.

Intestinal resorption of Ca^{2+}

$\text{Fc}_{240} = 9\%$ (normal limits 20 - 22 %)

Intestinal resorption of Ca^{2+}

$\text{Fc}_{240} = 9\%$ (normal limits 20 - 22 %)

Principles of examination

Resorption of Ca^{2+} is derived from Sr^{2+} resorption.
After p.o. administration of 2,5 mmol SrCl_2 , serum levels of Sr^{2+} are measured immediately (c_0) and after 240 min (c_{240}). Fc_{240} is calculated.

Causes of impaired gut resorption of Ca



Low intake of Ca in a food

Low biol. availability of Ca in a food

„Negative balance“ – secretion of Ca

Impaired absorption of Ca



Physiology of Ca absorption

**D-vitamin independent
paracelullular absorption**

D-vitamin dependent absorption



Ca²⁺ resorption is independent on a specific transport protein

Calcitriol changes conformation of **calmodulin**. More effective linkage of Ca²⁺ to calmodulin facilitates a passing of Ca²⁺ through intestinal wall

Calcitriol induces a synthesis of cytosolic **calbindin** (CaBP, calcium-binding protein). CaBP has high affinity to Ca²⁺ in cytosol. Therefore an increase of intracellular Ca²⁺ is limited and Ca is released from enterocytes.

Calcitriol induces a synthesis of **Ca²⁺-ATPase**, from Ca²⁺ transport from enterocyte do extracellular fluid.

Conclusion


Idiopathic impaired calcium intestinal resorption

- Recommended nutritional support by „sipping“.
- Therapy: Ca 1,5 g / day, Alpha-D3 1 ug / day.
- Follow up in out-patient department of osteology



Case report No. 3

Hypopituitarism



Female, 38 yr.

at age 20 umbilical herniotomy

at age 28 cholecystectomy for gallbladder stones

at age 34 sectio cesarea

Without regular medication

Abuse - negative

Allergy - negative

Social status: teacher, 3 children

Gynecology: 3 deliveries, 3 abortions

June 26, 2009, sectio cesarea as a 3rd delivery

One week after delivery, severe weakness, headache and impaired lateral vision revealed.

June 26, 2009, sectio cesarea as a 3rd delivery

One week after delivery, severe weakness, headache and impaired lateral vision revealed.



Perimeter

Bitemporal hemianopsia

MRI of pituitary (September 14, 2009)

Macroadenoma of pituitary gland with suprasellar propagation, in a near contact with chiasma n. optici.

Laboratory

Prolactin	117 mIU/l	(<630)
LH	0,62 IU/l	(0,7-5,0)
FSH	2,22 IU/l	(0,5-8,0)
Estradiol	11,0 pg/ml	(25-75)
free T4	6,2 pmol/l	(11,0-21,0)
TSH	0,6 mIU/l	(0,5-5,0)
Cortisol	81,0 nmol/l	(200-750)

All described hormones are borderline or **bellow normal limits**.

Laboratory findings respond to combined pituitary insufficiency:

- hypogonadism,
- hypothyroidism,
- hypocortisolism

Hospitalization at internal department

Substitution therapy of hypopituitarism

High doses of corticoids → hypokalemia

October 04, 2009 – paroxysm of ventricular fibrillation

Hospitalization at internal department

Substitution therapy of hypopituitarism

High doses of corticoids → hypokalemia

October 04, 2009 – paroxysm of ventricular fibrillation

**Transport to ICU of the 3rd Internal clinic, General
Faculty Hospital in Prague**

**Stabilization, correction of hypokalemia and
hypophosphatemia**

**Indicated to neurosurgical treatment of pituitary
adenoma**

Hospitalization at internal department

Substitution therapy of hypopituitarism

High doses of corticoids → hypokalemia

October 04, 2009 – paroxysm of ventricular fibrillation

**Transfer to ICU of the 3rd Internal clinic, General
Prague**

Hydrocortison 25-15-5 mg / d. hypokalemia and

Euthyrox 125 µg / d.

Omeprazol (Helicid) 20 mg / d. treatment of pituitary

January 25, 2010 – admission to Neurosurgical dept.

```
graph TD; A[January 25, 2010 – admission to Neurosurgical dept.] --> B[Perimeter Normal finding !!]; B --> C[MRI of pituitary (January 26, 2010, Prague) Hypophysis 12x6 mm, without signs of adenoma !! Infundibulum without dislocation. No signs of bleeding. Normal size of ventricular system.];
```

Perimeter

Normal finding !!

MRI of pituitary (January 26, 2010, Prague)

Hypophysis 12x6 mm, without signs of adenoma !!

Infundibulum without dislocation.

No signs of bleeding.

Normal size of ventricular system.

Conclusion:

Actual finding is not an indication to surgery.

Pituitary apoplexy ?

**Persisting panhypopituitarism, requiring
substitution.**

Following revision of previous MRI images



The finding, regarded to be a pituitary macroadenoma, is a typical image of lymphocyte (autoimmune) hypophysitis.

Development of MRI image as well as clinical course confirm this diagnosis.


Prof. J. Marek, 3rd Internal Dept., 1st Faculty of Medicine




Differential diagnostics of pituitary expansions




Differential diagnostics of pituitary expansions




35 % hormonally afunctional
30 % PRL ... prolactinoma
25 % STH ... acromegaly
10 % ACTH ... Cushing sy
< 1 % other (TSH, LH, FSH)



Differential diagnostics of pituitary expansions



Differential diagnostics of pituitary expansions




Autoimmune hypophysitis

The most frequent type of chronic pituitary inflammation

Prevalence: females – mainly during pregnancy + shortly after delivery

Frequent mismatch with macroadenoma


Good prognosis with tendency to spontaneous regression.




Patogenesis of hypophysitis

Hormonal dysbalance (gravidity, post-delivery)

„Dysbalance“ of T_{H1} / T_{H2} responses



Clinical findings



Local changes

47 % headache

33 % vision alteration

Hypopituitarism

33 % ↓ ACTH

13 % ↓ TSH

Hypothalamic - pituitary dysfunction

27 % diabetes insipidus

23 % ↑ PRL

Dif. dg. between hypophysitis and pituitary adenoma

Young females

Onset usually before or early after delivery

Early symptoms - headache, vision alteration

Spontaneous regression

Dif. dg. between hypophysitis and pituitary adenoma

MRI image ... elongated, pyriform shape of pituitary in hypophysitis



Dif. dg. between hypophysitis and pituitary adenoma

„Selective“ hypopituitarism:

Pituitary destruction caused by **tumor** leads usually to subsequent dysfunction in following order:

↓
LH, FSH
STH, TSH
↓
ACTH

In a case of lymphocyte **hypophysitis**, the decrease of ACTH is usually the first sign.